$$H = -\frac{h^2}{2m_N} \nabla_N^2 - \frac{h^2}{2m} \nabla_e^2 - \frac{KZe^2}{r}$$
nucleus electron — ①

We once again perform the CM. transform \tilde{r} \tilde{r}

14/11/20 Fec. 36.
The Halom Moken

briting the Schrödiger equalism in spherical polar coordinate we have:

$$-\frac{\hbar^2}{2\mu}\left[\frac{1}{7}\frac{\partial^2}{\partial r}\left(r\Psi(r,0,\varphi)\right) - \frac{1}{\hbar^2r^2}\left[\frac{1}{7}\Psi(r,0,\varphi)\right] - \frac{1}{\hbar^2r^2}\left[\frac{1}{7}\Psi(r,0,\varphi)\right] - \frac{1}{4^2r^2}\left[\frac{1}{7}\Psi(r,0,\varphi)\right] - \frac{1}{4^2r^2}\left[\frac{1}{7}\Psi(r,\varphi)\right] -$$

From (2) we expect that both R(r) and, hence, E could depend on I, but not on m. This means that each energy level will have a (x1+1) defencionly coming from i, evgen richs.

To simplify (7) Bet's consider

R(r) = U(r)/r - (8)

Substituting lies in & yields -V(r)
-, an effective
potential. -) Unlike the usual 1-1 egn. $, r = [0, \infty)$

$$V_{L}(r) = -R^{2}e^{-\frac{1}{2}} + \frac{L(RH)R^{2}}{2\mu r^{2}}$$

repulsive centrifugal

Larrier for $L>0$.

Boundary conditions on $U(r)$ are

cultivest from a regular 1-d

problem

(and be rewritten as an eigenvalue equ.

$$\hat{D}_{L}U(r) = EU(r) - [u]$$

when
$$\hat{D}_{L} = \begin{bmatrix} -\frac{1}{2}A^{2} & +V(r) \\ \frac{1}{2}H & dr^{2} & +V(r) \end{bmatrix} - [2]$$

Since \hat{D}_{e} has to be Hermitian we have,

for 2 functions $U_{l}(r) = \frac{1}{2}U_{l}(r) = \frac{1}{2}U_{l}(r)$

That is.
$$\int_{0}^{\infty} U_{1}^{*} \left(\hat{D}_{\ell} U_{2} \right) dr = \left(\int_{0}^{\infty} U_{2}^{*} \left(\hat{D}_{\ell} U_{1} \right) dr \right)^{*}$$

$$= \int_{0}^{\infty} \left(\hat{D}_{\ell} U_{1} \right)^{*} U_{2} dr \quad (14)$$

$$=\int_{0}^{\infty} \left(\hat{D}_{2} V_{1} \right)^{*} V_{2} dr - 19$$

 $\int dr \left[U_{1}^{*} \left(-\frac{h^{2}}{2\mu} \frac{d^{2}}{2r^{2}} \right) U_{2} \right] = 0$ $- U_{2} \left(-\frac{h^{2}}{2\mu} \frac{d^{2}}{J_{12}} \right) = 0$ - (15)

Since V (1) is a multiplication or space-local operator).

Inlégrating each lern in (15) by parts we jet $\left(\begin{array}{cccc}
U_1^* & \underline{d} & U_2 & - & U_2 & \underline{d} & U_1^* \\
d & \overline{d} & \overline{r} & - & \overline{d} & \overline{r}
\end{array}\right) = 0$

Therefore, solution to @ must satisfy (6). Normalischer Pg. 4, for bound steller, $\int \int \int |\psi(r,0,+)|^2 dr \sin \theta d\theta d\theta$ $= \int (P(r))^2 r^2 dr \int d\Omega |\chi(\Omega)|^2$ = 1

 $\int_{0}^{\infty} |\mathcal{L}(r)| r^{2} d\tau = 1 \quad -(3)$

or $\int_{0}^{\infty} \left| \left| v(r) \right|^{2} dr = 1 \qquad -(8)$ That is for bound states

 $\mu \to 0$ $\mu \to 0$

For either case,

$$\begin{vmatrix} u_1^* & dv_2 & -v_2 & dv_1^* \\ dv & dv \end{vmatrix} = 0$$
 $\begin{vmatrix} u_1^* & dv_2 & -v_2 & dv_1^* \\ dv & dv \end{vmatrix} = 0$
 $\begin{vmatrix} v_1^* & dv_2 & -v_2 & dv_1^* \\ dv & dv \end{vmatrix} = 0$

en this case

$$\Rightarrow \text{ for } l=0, m=0 \qquad \forall n \subseteq Y_0^0 - 25$$
and
$$\nabla^2 \left(\frac{1}{7}\right) = -478 (\bar{r}) \qquad -25$$

 $\nabla \left(\frac{1}{7}\right) = -4110 (F)$ \hat{U} it doesnit salisfig llie Schröduzer

egn. at llie ongin

e we must have that c=0 or $V \longrightarrow 0 \longrightarrow 0$

Now as r-so, the dominant lein in the potential is the centripyed lein is l(2+1) >> 1 and >> E.

 $\frac{d^2U}{d\tau^2} = \frac{l(l+1)}{r^2} U - \frac{29}{r^2}$

A brial solution of the form

U(r) ~ r yields $\alpha(\alpha-1)$ $\gamma^{\alpha-2} = \lambda(\alpha+1)$. γ^{α} & a=l+1 & -l. -(31) doent derent

(doent derent

on E)

Thous up at T-0 against our

rondition (27). So we reject it. :. U(r) ~ rl+1 (r→0) (33)

Consider now the behavior of U as T-x0.

V(x) -> 0 as T-x0 in our case.

At large the qu (3) becomes.

(3b) =)
$$U_{E}(r\rightarrow a)$$
 $N \in \mathbb{R}^{p}$ (3b) where $K_{E} = \frac{2\mu IE}{R^{2}}$ (3b) There are some subtleties to this expectation coming from the $1/r$ decay of the potential But motivated by this behaviour, we can define $U(r) = f(r) \in \mathbb{R}^{p}$ (3f)

Then
$$V_{ER}(P) = e^{-\beta} f_{ER}(P) - 4D$$
 f then satisfie:

$$\frac{d^2f}{dP} - 2\frac{df}{dP} + \left[\frac{KZe^2\lambda}{P} - \frac{L(l+1)}{P^2}\right]f$$

where $\lambda = \sqrt{\frac{2m}{k^2|E|}} - 43$

We by a phynomial expansion for f motivated by this $(-\infty)$ limit behaviour love channel earlier. Since $U(P) \sim P^{l+1}(P \rightarrow \infty)$ we can shork $f(P) = P^{l+1}(P \rightarrow \infty)$ we can shork $f(P) = P^{l+1}(P \rightarrow \infty)$ we can shork $f(P) = P^{l+1}(P \rightarrow \infty)$ and

collectuy same power leven Toythe we get $-(2Ke^2)\lambda + 2(k+l+l)$ (k+l+2) (k+l+1) -l(l+1) 4 Energy levels: Now, the Cent = 2 k -(45)

This is the same behaviour of coeff, we would get for the power sens,

Of VIG. pme^{2P}. 46)

So we are tempted to equale

f = V - 47

But, this would imply

$$U = \int_{0}^{m} e^{2\rho} \cdot e^{-\rho} = \int_{0}^{m} e^{\rho} - 48$$

So we must terminale the series at some finite k . This would happen of, for some k , we have $C_{kH} = 0$ -49

$$= \frac{-49}{c_k} = 0 \qquad -49$$

$$= \frac{1}{c_k} = 0 \qquad -49$$

or, from definition of
$$X$$
:
$$E = -|E| \qquad (E < 5)$$

$$= \frac{2m}{h^2} / \lambda^2$$

$$= \frac{-m / (x^2 e^2)^2}{2 + max^2 (k+l+l)^2}$$
Thus, is not fixed here beyond

Since kman is not fixed here beyond

being a non-negative integer

we get energy quantization

for
$$k_{max} = 0, 1, 2, \cdots$$
 $k_{max} = 0, 1, 2, \cdots$
 $k_{max} = 0, 1$

$$\Rightarrow E_n = -\frac{Z^2}{n^2} R_y - 64$$

1 hy a 13.6 eV.

Wavefunctions: lyven lie recursion

Prelations and the kmax = n-1-1

we can

work batk the wavefor.

The corresponding of are a set of polynomials called lie Associate Lagueur polynomials dendid by

where
$$L_p(x) = (-1)^k \frac{d^k}{dz^k} L_{p+k} - \frac{\pi}{2}$$

$$L_p(x) = e^{x} \left(\frac{d^k}{dx^p}\right) \left(e^{-x}x^p\right) + \frac{\pi}{2}$$
with this definition we get
$$R_{nk}(x) \sim e^{-x/na_0} \left(\frac{\gamma}{na_0}\right)^{k} L_{n-k-1}(\frac{2r}{na_0})$$

$$-\frac{\pi}{2}$$

Allowed values of l for a given on are restricted by the highest degree allowed for L

ie l = n-1, n-2, -3, 2, 1, 0 $f \neq p \leq \frac{1}{2}$

$$\psi_{100}(\bar{r}) = \frac{1}{123} \frac{1}{2} e^{-T/ab} - E$$

$$\psi_{20}(\bar{r}) = \frac{1}{32\pi a_0^3} \frac{1}{2} e^{-T/2ab} - E$$

$$\psi_{210}(\bar{r}) = \frac{1}{(32\pi a_0^3)} \frac{7}{ab} e^{-T/2ab} e^{-T/2ab}$$

$$\psi_{210}(\bar{r}) = \frac{1}{(32\pi a_0^3)} \frac{7}{ab} e^{-T/2ab}$$

$$\psi_{210}(\bar{r}) = \frac{1}{(32\pi a_0^3)} \frac{7}{ab}$$

$$\psi_{21$$

The normalized légenfundion of ller Halois au llier:

$$\frac{4^{2}py}{2^{1}} = \frac{4^{2}11}{2^{1}} + \frac{4^{2}1-1}{2^{1}}$$

$$= -\left(\frac{1}{64\pi a_{0}^{3}}\right) \frac{r}{a_{0}} e^{-\frac{r}{2}a_{0}}$$

$$= -\left(6\right) \frac{y}{5}$$
Similar combinations are possible

for d'orbital.


Note Mat Me Real orbitals

are NOT eigenfunctions of Lz

but continue to be eigenfunctions of

12

Plots of the radial Junchen



Source: MIT Open courseware

Probability of finding the electrons in a spherical shell of radius of and thetenen dr. $= \int_{r}^{r+r} |\Psi(r)|^{2} dr d\Omega$ $= \int_{r}^{r} P(r) r^{2} dr$ In the ground (15) state this $\alpha e^{-2r/a_0} r^2 dr$ feates a maximum when


d (e st/ao r)

d T Hence, the Bohr radius has a physical meaning. Plots of orbitals.

There are 3-d functions so hard to plot the the sold as isosurfaces of density. That is an image shows the surface made by points (F, 4) having the same density |4|2.

A nice visualization of the orbitals can be found at .. This Site

A rough sketch of the spatial orientations of the orbitals are given below.



The shaded unshaded regions denote that the wavefuries is positive, negative.

At the boundary of the shaled & unshaded regions (pts, line, planes, et) the wavefunction goes to zero yielding a node. Unem(r,o, φ) ~ Rne(r) P, (ωσ) eing 7 L 28+1 (2Kr)

degræ n-l-1 Segree l n-l-1 Radiaf

l angulat nodes